



Does Glucose Variability Influence the Relationship Between Mean Plasma Glucose and HbA_{1c} Levels in Type 1 and Type 2 Diabetic Patients?

Citation

Kuenen, Judith C., Rikke Borg, Dirk J. Kuik, Hui Zheng, David Schoenfeld, Michaela Diamant, David M. Nathan, and Robert J. Heine. 2011. Does glucose variability influence the relationship between mean plasma glucose and HbA_{1c} levels in type 1 and type 2 diabetic patients?. *Diabetes Care* 34(8): 1843-1847.

Published Version

doi:10.2337/dc10-2217

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10464973>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Does Glucose Variability Influence the Relationship Between Mean Plasma Glucose and HbA_{1c} Levels in Type 1 and Type 2 Diabetic Patients?

JUDITH C. KUENEN, MD¹
RIKKE BORG, MD²
DIRK J. KUIK, MSC³
HUI ZHENG, PHD⁴
DAVID SCHOENFELD, PHD⁴

MICHAELA DIAMANT, MD¹
DAVID M. NATHAN, MD⁵
ROBERT J. HEINE, MD⁶
ON BEHALF OF THE ADAG STUDY GROUP*

OBJECTIVE—The A1C-Derived Average Glucose (ADAG) study demonstrated a linear relationship between HbA_{1c} and mean plasma glucose (MPG). As glucose variability (GV) may contribute to glycation, we examined the association of several glucose variability indices and the MPG-HbA_{1c} relationship.

RESEARCH DESIGN AND METHODS—Analyses included 268 patients with type 1 diabetes and 159 with type 2 diabetes. MPG during 3 months was calculated from 7-point self-monitored plasma glucose and continuous glucose monitoring. We calculated three different measures of GV and used a multiple-step regression model to determine the contribution of the respective GV measures to the MPG-HbA_{1c} relationship.

RESULTS—GV, as reflected by SD and continuous overlapping net glycemic action, had a significant effect on the MPG-HbA_{1c} relationship in type 1 diabetic patients so that high GV led to a higher HbA_{1c} level for the same MPG. In type 1 diabetes, the impact of confounding and effect modification of a low versus high SD at an MPG level of 160 mg/dL on the HbA_{1c} level is 7.02 vs. 7.43 and 6.96 vs. 7.41. All GV measures showed the same tendency.

CONCLUSIONS—In only type 1 diabetic patients, GV shows a significant interaction with MPG in the association with HbA_{1c}. This effect is more pronounced at higher HbA_{1c} levels. However, the impact of GV on the HbA_{1c} level in type 1 diabetes is modest, particularly when HbA_{1c} is close to the treatment target of 7%.

Diabetes Care 34:1843–1847, 2011

Since the Diabetes Control and Complications Trial (DCCT) and the UK Prospective Diabetes Study (UKPDS) (1,2) established the relationship between HbA_{1c} and the development of long-term diabetes complications, HbA_{1c} has become the key monitoring tool in diabetes management.

During the lifetime of the erythrocyte, hemoglobin (Hb) is gradually glycated. The proportion of the glycated sites, HbA_{1c}, within the erythrocyte increases throughout its life span and reflect the exposure to mean blood glucose (MBG) levels during the preceding 2–3 months (3). This non-enzymatic posttranslational modification is

relatively slow. In vivo and in vitro studies have shown that HbA_{1c} levels are directly proportional to the time-averaged concentration of glucose during the erythrocyte's life span (3–6). Given the kinetics of glycation, brief periods of hyperglycemia should not have a major impact on HbA_{1c} levels (7–9).

However, increased glycated protein levels are documented in some nondiabetic pathological states. So, hyperglycemia is not the complete answer to the etiology of increased early glycated products in nondiabetic conditions. A common denominator is oxidative stress. It has been hypothesized that oxidative stress either via increasing reactive oxygen species or by depleting the antioxidants may modulate the genesis of early glycated proteins in vivo (10,11). Hyperglycemia stimulates oxidative stress (12) and glucose variability; in particular, postprandial glucose excursions have been regarded as potentially deleterious as a result of, among other factors, their association with the increase of oxidative stress (13). Therefore, glucose variability (GV) could influence the glycation of HbA_{1c}.

Previous studies have examined whether the relationship between mean plasma glucose (MPG) levels and HbA_{1c} is influenced by glucose variability and found no or minimal influence (10,14,15). However, these studies used limited self-monitoring of blood glucose (SMBG) data to assess mean glucose levels and variability in relatively small numbers of measurements. These methods could underestimate glycemic excursions. Continuous glucose monitoring (CGM) provides a more complete view of glycemic excursions, including the duration and frequency of the excursions, and allows calculation of features of GV. Our aim was to examine the influence of GV on the MPG-HbA_{1c} relationship in the A1C-Derived Average Glucose (ADAG) study.

RESEARCH DESIGN AND METHODS

The ADAG study was conducted at 10 centers in the U.S.,

From the ¹Diabetes Center, VU University Medical Center, Amsterdam, the Netherlands; the ²Steno Diabetes Center, Copenhagen, Denmark; the ³Department of Epidemiology and Biostatistics, VU University Medical Center, Amsterdam, the Netherlands; the ⁴Biostatistics Center, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts; the ⁵Diabetes Center, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts; and ⁶Lilly, Minneapolis, Minnesota.

Corresponding author: Judith C. Kuenen, jc.kuenen@me.com.

Received 2 December 2010 and accepted 12 May 2011.

DOI: 10.2337/dc10-2217

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc10-2217/-/DC1>.

*A complete list of the members of the ADAG Study Group can be found in the Supplementary Data.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Europe, and Africa from 2006 to 2008 to define the relationship between HbA_{1c} and average glucose levels. Because a full description of this observational study has been published (14), we describe it here only briefly. A total of 268 individuals with type 1 diabetes and 159 individuals with type 2 diabetes (age 18–70 years) completed the study. Participants were selected based on stable glycemic control as evidenced by two HbA_{1c} values within one percentage point of each other in the 6 months prior to recruitment. Individuals with a wide range of HbA_{1c} levels were included.

Participants with conditions leading to major changes in glycemia (infectious disease, steroid therapy, and pregnancy) or conditions that might interfere with the measurement of HbA_{1c} or the relationship between HbA_{1c} and MPG (hemoglobinopathies [16], anemia, increased erythrocyte turnover, blood loss and/or transfusions, or chronic renal or liver disease) were excluded (14). The study was approved by the human studies committees at the participating institutions, and informed consent was obtained from all participants.

Measurements of glycemia

During the study period, CGM (Medtronic Minimed, Northridge, CA) was performed at home four times with 4-week intervals during the 16-week study period. Monitoring period lasted at least 48 h, during which time glucose levels were assessed every 5 min. CGM data were accepted for analysis if there were no gaps longer than 120 min and if the mean absolute difference with the Hemocue calibration results was <18%, as recommended by the manufacturer. For calibration purposes, participants performed SMBG with the Hemocue meter (Hemocue Glucose 201 plus; Hemocue, Ångelholm, Sweden) during the days of CGM.

For adequate calculation of MPG, subjects additionally performed a seven-point SMBG (OneTouch Ultra; Lifescan, Milpitas, CA) for at least 3 days per week during the weeks when CGM was not performed. All blood glucose values stated are plasma equivalents.

HbA_{1c} samples were analyzed with four highly intercorrelated DCCT-aligned assays: a high-performance liquid chromatography assay, two immunoassays, and an affinity assay (all approved by the National Glycohemoglobin Study Program). The mean HbA_{1c} value at the end of the 12 week study period was used (14).

Calculating glucose variability

Three indices of intraday glucose variability were calculated based on CGM: the SD of mean glucose concentrations, the mean amplitude of glycemic excursions (MAGE), and the continuous overlapping net glycemic action (CONGA). High SD, MAGE, and CONGA values indicate high intraday glucose variability. MAGE is the mean of the differences between consecutive peaks and nadirs, only including changes of >1 SD of glycemic values and thus capturing only major fluctuations (17). For the calculation of CONGA_n, the difference of the current value compared with the value *n* hours previously was calculated for each observation after the first *n* hours. The CONGA_n is the SD of these differences (18). In the analyses, we used CONGA at 4 h (CONGA₄). Calculations based on CGM data were calculated after exclusion of the initial 2 h of monitoring, which is considered to be an unstable calibration period.

Statistical analysis

First, we explored the correlations between MPG and HbA_{1c} and measures of glycemic variability as SD, MAGE, and/or CONGA₄ for the total diabetic population and the two diabetes types. Multiple linear regression was used to investigate confounding and effect-modifying influence of clinical parameters (glycemic variability) on the relation between the determinant (MPG) and outcome (HbA_{1c}) of interest. We then assessed which of the variability measures (SD, MAGE, or CONGA₄) had the strongest impact on the MPG-HbA_{1c} relationship by confounding or effect modification.

Effect modification was concluded when the slope of the interaction term of glycemic variability and determinant was significant. If no effect modification might be concluded, a parameter ΔB was computed as the relative difference of the slope of the determinant in the model without and with the clinical parameter. Confounding was concluded when the absolute value of ΔB exceeded the generally accepted threshold of 10%.

Multivariate confounding was investigated with a variant of stepwise regression, in which the stepping criterion was not a *P* value but the ΔB as long as it exceeded the threshold. For significance, a threshold of $\alpha = 0.05$ was used.

Analyses were done for the total population and stratified for the type of diabetes. Finally, we illustrated the magnitude of the effect caused by the variability indices,

by confounding or effect modification, on the MPG-HbA_{1c} relationship.

RESULTS—Of the 507 patients enrolled, 427 completed the study and had adequate glucose monitoring and HbA_{1c} samples for inclusion in the analyses. Two hundred and sixty-eight participants had type 1 diabetes, and 159 had type 2 diabetes. The CGM and the SMBG data during the 3-month period included approximately 2,400 and 300 measurements per subject, respectively. The relationship between the HbA_{1c} level at the end of the 3-month study period and MPG calculated over the preceding 3 months was expressed as the simple linear regressions. The formula for the total diabetic population was as follows: HbA_{1c} (%) = $0.028 \times \text{MPG (mg/dL)} + 2.66$ ($R^2 = 0.80$). The formula for type 1 diabetes was as follows: HbA_{1c} (%) = $0.028 \times \text{MPG (mg/dL)} + 2.77$ ($R^2 = 0.77$). The formula for type 2 diabetes was as follows: HbA_{1c} (%) = $0.028 \times \text{MPG (mg/dL)} + 2.62$ ($R^2 = 0.82$).

The clinical and glycemic characteristics are shown in Table 1. Mean HbA_{1c} (SD) for type 1 diabetic patients was 7.3% (1.1) and for type 2 diabetic patients was 6.8% (1.1).

All GV measures had significant influence on the MPG-HbA_{1c} relationship for the total population. The variability index SD showed the strongest influence on the MPG-HbA_{1c} relationship. None of the GV measures showed confounding for all diabetic patients pooled or for the type 1 or type 2 diabetic patients separately (Table 2).

In the type 1 diabetic patients, the effect modification of SD and CONGA₄ was significant ($P < 0.01$ and $P = 0.02$), and for the MAGE it was just not significant ($P = 0.06$) (Table 2). The MPG/HbA_{1c} linear regression formula with confounding for type 2 diabetes was as follows: HbA_{1c} (%) = $2.64 + 2.63 \times \text{MPG}/100 + 0.58 \times \text{SD}/100$. The MPG-HbA_{1c} linear regression formula with effect modification for type 1 diabetes was as follows: HbA_{1c} (%) = $3.91 + 1.79 \times \text{MPG}/100 - 1.37 \times \text{SD}/100 + 1.25 \times \text{MPG}/100 \times \text{SD}/100$. The impact of effect modification of low GV (SD = 30 mg/dL) versus high GV (SD = 100 mg/dL) for an MPG level of 160 mg/dL in type 1 diabetes on the HbA_{1c} level was 6.96 vs. 7.41%, respectively, as shown in Table 3. At an MPG level of 220 mg/dL (HbA_{1c} following the regression formula of 8.89%), a decline in the SD parameter from 100 to 30 mg/dL reduced HbA_{1c} from 9.23 to 8.26%.

Table 1—Baseline clinical and glycemic characteristics

	All	Type 1 diabetes	Type 2 diabetes
n	427	268	159
Age (years)	47.6 ± 13.6	44.1 ± 12.9	56.6 ± 9.4
Sex (% female)	54	52	51
Ethnicity (% non-Hispanic whites)	83	93	73
Current smokers	11	12	9
Insulin treatment	76	100	38
Glycemic measures			
HbA _{1c} (%)	6.8 ± 1.3	7.3 ± 1.1	6.8 ± 1.1
MPG (mg/dL)	149.4 ± 39.6	162 ± 36	149.4 ± 36
Measures of GV			
CGM SD (mg/dL)	48.6 ± 25.2	64.8 ± 16.2	39.6 ± 16.2
MAGE (mg/dL)	86.4 ± 43.2	115.2 ± 32.4	68.4 ± 27
CONGA ₄ (mg/dL)	66.6 ± 28.8	88.2 ± 23.4	52.2 ± 21.6
SD (mg/dL)			
≤30	61 (14.3)	9 (3.4)	52 (32.7)
<30–60	173 (40.5)	84 (31.3)	89 (56)
<60–90	173 (40.5)	155 (57.8)	18 (11.3)
>90	20 (4.7)	20 (7.5)	0 (0)

Data are means ± SD, %, or n (%).

For all patients pooled, there was no effect modification of the respective GV measures on the MPG-HbA_{1c} relationship. For type 2 diabetic patients, the impact of effect modification from the respective GV measures was far from significant (Table 2). The number of patients with a predefined SD is shown in Table 1 for all patients pooled and for the type 1 and type 2 diabetic patients separately.

CONCLUSIONS—This study demonstrated a significant effect of GV, as reflected by SD, on the MPG-HbA_{1c} relationship. High GV (SD) is associated with

higher HbA_{1c} levels for a given MPG, and this effect was more pronounced at higher HbA_{1c} and MPG values. However, the magnitude of this effect of GV was small and only demonstrable in type 1 diabetic patients. Possibly, the type 2 diabetic patient group was too small (*n* = 159) and the variability in this group too low to find this interaction.

The ADAG study showed a tight correlation between HbA_{1c} and MPG, allowing the translation of HbA_{1c} into estimated average glucose (14,19). It has been suggested that GV could affect the MPG-HbA_{1c} relationship, but this has not

previously been demonstrated (20–22). To our knowledge, the current study is the largest study reporting an influence of GV—as expressed by SD, MAGE, and CONGA₄ calculated from CGM—on the MPG-HbA_{1c} relationship. The discrepancies in the MPG-HbA_{1c} relationship are less likely caused by technical errors because this study included accurate and centralized measurements of HbA_{1c} values and intensively measured plasma glucose concentrations (~2,700 values) in a large and diverse population. Also, individuals with conditions or treatment that might result in major changes in glycemia or interference with the HbA_{1c} assay, or the MPG-HbA_{1c} relationship, were excluded. These precautions allowed us to search for factors other than MPG that may contribute to HbA_{1c}.

In general, GV is higher in patients with poor glycemic control and in type 1 diabetic patients compared with type 2 diabetic patients, which can be attributed to insulin therapy and higher insulin sensitivity. High GV may affect glycation because of periodic exposure of the erythrocyte to high glucose levels and therefore to faster irreversible glycation.

Other factors like hyperglycemia-induced oxidative stress may affect the glycation process. In recent literature, it has been speculated that oxygen free radicals per se or with an associated decrease in antioxidants may modulate the formation of early glycated protein (10,11).

Brownlee (12) demonstrated that hyperglycemia stimulates oxidative stress. High GV and especially postprandial glucose excursions were also previously

Table 2—The P values of the influence of the respective GV measures themselves, as well as effect modification and the Δ of confounding, calculated from the respective slopes (B and B') from the regression equations, on the HbA_{1c}-MPG relationship for all patients pooled and for type 1 and type 2 diabetic patients separately

	Influence of the GV measure (P)	Slope of MPG (B) in the main regression formula	Slope of MPG (B') in the regression formula with the GV measure	ΔConfounding (%)*	Effect modification (P)
SD					
All	<0.01	2.818	2.624	6.9	0.06
Type 1 diabetic	0.01	2.781	2.631	5.4	<0.01
Type 2 diabetic	0.06	2.782	2.637	5.2	0.74
MAGE					
All	<0.01	2.818	2.700	4.2	0.37
Type 2 diabetic	0.19	2.781	2.721	2.2	0.06
Type 2 diabetic	0.19	2.782	2.698	3.0	0.19
CONGA ₄					
All	<0.01	2.818	2.667	5.4	0.15
Type 1 diabetic	0.06	2.781	2.687	3.4	0.02
Type 2 diabetic	0.07	2.782	2.661	4.3	0.46

*ΔConfounding in % = 100 × absolute (B' – B)/B.

Table 3—Quantification or impact of confounding or effect modification for type 1 diabetic patients of a low vs. a high SD for a given MPG on the HbA_{1c} level next to the HbA_{1c} values calculated with the regression formula for type 1 diabetes

MPG [mg/dL (mmol/L)]	Regression formula	HbA _{1c} (%)			
		Confounding		Effect modification	
		SD 30 mg/dL	SD 100 mg/dL	SD 30 mg/dL	SD 100 mg/dL
140 (7.8)	6.67	6.50	6.90	6.53	6.80
160 (8.9)	7.22	7.02	7.43	6.96	7.41
180 (10)	7.78	7.55	7.95	7.39	8.02
200 (11.1)	8.34	8.08	8.48	7.83	8.62
220 (12.2)	8.89	8.60	9.01	8.26	9.23
240 (13.3)	9.45	9.13	9.53	8.69	9.84

associated with oxidative stress in type 2 diabetes (13). The activation of oxidative stress, estimated from urinary excretion rates of isoprostanes, was highly correlated with MAGE calculated from CGM (13). However, Wentholt et al. (23) could not replicate these results in type 1 diabetes. Recently, Ceriello et al. (15) demonstrated that high intraday GV was more damaging to endothelial function than stable hyperglycemia and that oxidative stress influences glycation needs to be determined.

On the other hand, it has been demonstrated that erythrocyte survival is shorter at chronic high glucose concentrations levels, which might falsely lower HbA_{1c} levels. Peterson et al. (24) showed that the life span of ⁵¹Cr-labeled erythrocytes increased in all seven subjects when their poorly controlled diabetes was adequately treated. Virtue et al. (25) concluded that there is a hyperglycemia-related decrease in erythrocyte survival as measured by carbon monoxide in the expired air, which results in an exponential underestimation of the severity of hyperglycemia at higher HbA_{1c} levels. Similarly, hyperglycemia-related osmotic stress may influence erythrocyte permeability and could cause damage to the erythrocyte and shorten its life span. These findings could lead to underestimation of HbA_{1c} at higher MPG levels, concealing a glycemic control worse than indicated by HbA_{1c} measurements. However, we found that type 1 diabetic patients with high GV display higher HbA_{1c} levels than suspected based on MPG. This effect was more pronounced at higher HbA_{1c} levels, indicating that focus on reducing GV, especially in patients with poor glycemic control, could help reduce HbA_{1c} levels.

Limitations of our study are that CGM has a limited range of reliable measurements between 2.2 mmol/L and 22.2 mmol/L. Therefore, theoretically, CGM performance could be less precise in patients with high glycemic variability. Furthermore, CGM has a lag time in glucose values compared with the venous measured values (the physiological gap), and this can result in underestimation of the influence of GV on the glycation of HbA_{1c}, and no measures of erythrocyte survival, oxidative stress, or clinical follow-up are available in this population.

In conclusion, at higher levels of GV the relationship between HbA_{1c} and MPG in patients with type 1 diabetes is altered, leading to a higher HbA_{1c} level for a given MPG. However, the impact (near the HbA_{1c} treatment target of 7%) is only modest.

Acknowledgments—The ADAG study is supported by research grants from the American Diabetes Association and the European Association for the Study of Diabetes.

Financial support was provided by Abbott Diabetes Care, Bayer Healthcare, GlaxoSmithKline, sanofi-aventis Netherlands, Merck & Company, LifeScan, and Medtronic Minimed. Supplies and equipment were provided by Medtronic Minimed, LifeScan, and Hemocue. R.J.H. is employed by and owns stocks of Eli Lilly and Company. No other potential conflicts of interest relevant to this article were reported.

J.C.K. researched data, contributed to discussion, and wrote the manuscript. R.B. researched data, contributed to discussion, and edited the manuscript. D.J.K., H.Z., and D.S. researched data. M.D. contributed to discussion and edited the manuscript. D.M.N. reviewed and edited the manuscript. R.J.H. contributed to discussion and reviewed and edited the manuscript.

Parts of this study were presented in abstract form at the 44th Annual Meeting of the European Association for the Study of Diabetes

(Poster number 1065), 7–11 September 2008, Rome, Italy and at the 68th Scientific Sessions of the American Diabetes Association, San Francisco, California, 6–10 June 2008.

References

1. The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977–986
2. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–853
3. Bunn HF, Haney DN, Kamin S, Gabbay KH, Gallop PM. The biosynthesis of human hemoglobin A_{1c}. Slow glycosylation of hemoglobin in vivo. *J Clin Invest* 1976; 57:1652–1659
4. Beach KW. A theoretical model to predict the behavior of glycosylated hemoglobin levels. *J Theor Biol* 1979;81:547–561
5. Flückiger R, Winterhalter KH. In vitro synthesis of hemoglobin A_{1c}. *FEBS Lett* 1976;71:356–360
6. Higgins PJ, Bunn HF. Kinetic analysis of the nonenzymatic glycosylation of hemoglobin. *J Biol Chem* 1981;256:5204–5208
7. Bunn HF, Gabbay KH, Gallop PM. The glycosylation of hemoglobin: relevance to diabetes mellitus. *Science* 1978;200:21–27
8. Gonen B, Rubenstein A, Rochman H, Tanega SP, Horwitz DL. Haemoglobin A_{1c}: an indicator of the metabolic control of diabetic patients. *Lancet* 1977;2:734–737
9. Koenig RJ, Peterson CM, Jones RL, Saudek C, Lehrman M, Cerami A. Correlation of glucose regulation and hemoglobin A_{1c} in diabetes mellitus. *N Engl J Med* 1976;295:417–420
10. Selvaraj N, Bobby Z, Sathiyapriya V. Effect of lipid peroxides and antioxidants on glycation of hemoglobin: an in vitro study on human erythrocytes. *Clin Chim Acta* 2006;366:190–195
11. Selvaraj N, Bobby Z, Sridhar MG. Oxidative stress: does it play a role in the genesis of early glycosylated proteins? *Med Hypotheses* 2008;70:265–268
12. Brownlee M. A radical explanation for glucose-induced beta cell dysfunction. *J Clin Invest* 2003;112:1788–1790
13. Monnier L, Mas E, Ginot C, et al. Activation of oxidative stress by acute glucose fluctuations compared with sustained chronic hyperglycemia in patients with type 2 diabetes. *JAMA* 2006;295:1681–1687
14. Nathan DM, Kuenen J, Borg R, Zheng H, Schoenfeld D, Heine RJ; A_{1c}-Derived Average Glucose Study Group. Translating

- the A1C assay into estimated average glucose values. *Diabetes Care* 2008;31:1473–1478
15. Ceriello A, Esposito K, Piconi L, et al. Oscillating glucose is more deleterious to endothelial function and oxidative stress than mean glucose in normal and type 2 diabetic patients. *Diabetes* 2008;57:1349–1354
 16. Bry L, Chen PC, Sacks DB. Effects of hemoglobin variants and chemically modified derivatives on assays for glycohemoglobin. *Clin Chem* 2001;47:153–163
 17. Service FJ, Molnar GD, Rosevear JW, Ackerman E, Gatewood LC, Taylor WF. Mean amplitude of glycemic excursions, a measure of diabetic instability. *Diabetes* 1970;19:644–655
 18. McDonnell CM, Donath SM, Vidmar SI, Werther GA, Cameron FJ. A novel approach to continuous glucose analysis utilizing glycemic variation. *Diabetes Technol Ther* 2005;7:253–263
 19. Consensus Committee. Consensus statement on the Worldwide Standardization of the Hemoglobin A1C Measurement: the American Diabetes Association, European Association for the Study of Diabetes, International Federation of Clinical Chemistry and Laboratory Medicine, and the International Diabetes Federation. *Diabetes Care* 2007;30:2399–2400
 20. Derr R, Garrett E, Stacy GA, Saudek CD. Is HbA(1c) affected by glycemic instability? *Diabetes Care* 2003;26:2728–2733
 21. Service FJ, O'Brien PC. Influence of glycemic variables on hemoglobin A1c. *Endocr Pract* 2007;13:350–354
 22. McCarter RJ, Hempe JM, Chalew SA. Mean blood glucose and biological variation have greater influence on HbA1c levels than glucose instability: an analysis of data from the Diabetes Control and Complications Trial. *Diabetes Care* 2006;29:352–355
 23. Wentholt IM, Kulik W, Michels RP, Hoekstra JB, DeVries JH. Glucose fluctuations and activation of oxidative stress in patients with type 1 diabetes. *Diabetologia* 2008;51:183–190
 24. Peterson CM, Jones RL, Koenig RJ, Melvin ET, Lehrman ML. Reversible hematologic sequelae of diabetes mellitus. *Ann Intern Med* 1977;86:425–429
 25. Virtue MA, Furne JK, Nuttall FQ, Levitt MD. Relationship between GHB concentration and erythrocyte survival determined from breath carbon monoxide concentration. *Diabetes Care* 2004;27:931–935